

A FIRST-COUNTABLE NON-REMAINDER OF $\mathbb{H}$

ALAN DOW† AND KLAAS PIETER HART

ABSTRACT. We give a (consistent) example of a first-countable continuum that is not a remainder of the real line.

INTRODUCTION

The purpose of this note is to confirm a suspicion raised in [55; 5, Question 4.2]: we show that Bell's example, from [33, 3], of a first-countable compact space that is not an $\mathbb{N}^*$ -image can be adapted to produce admits a connected variation that is neither an $\mathbb{N}^*$ -image nor an $\mathbb{H}^*$ -image. The interest in this variation stems from the authors' version of Parovičenko's theorem from [99, 9]. That theorem states that every compact Hausdorff space of weight $\aleph_1$ or less is an $\mathbb{N}^*$ -image; the Continuum Hypothesis then implies that the $\mathbb{N}^*$ -images are exactly the compact Hausdorff spaces of weight $2^{\aleph_0}$ or less. We proved in [44, 4] a parallel result for $\mathbb{H}^*$ and continua (connected compact Hausdorff spaces). Since, by Arkhangel'skii's theorem [11, 1], first-countable compact spaces have weight at most $2^{\aleph_0}$ it follows that under CH first-countable compacta/continua are $\mathbb{N}^*$ -images/ $\mathbb{H}^*$ -images respectively.

Bell's graph. A major ingredient in our construction is Bell's graph, constructed in [22, 2]. It is a graph on the ordinal ω_2 , represented by a symmetric subset E of ω_2^2 . The crucial property of this graph is that there is *no* map $\varphi : \omega_2 \rightarrow \mathcal{P}(\mathbb{N})$ that represents this graph in the sense that $\langle \alpha, \beta \rangle \in E$ if and only if $\varphi(\alpha) \cap \varphi(\beta)$ is infinite.

Bell's graph exists in any forcing extension in which $\aleph_2$ Cohen reals are added; for the reader's convenience we shall describe the construction of E and adapt Bell's proof so that it applies to continuous maps defined on $\mathbb{H}^*$.

A FIRST-COUNTABLE CONTINUUM

Our starting point is a connected version of the Alexandroff double of the unit interval. We topologize the unit square as follows.

- (1) a local base at points of the form $\langle x, 0 \rangle$ consists of the sets

$$U(x, 0, n) = (x - 2^{-n}, x + 2^{-n}) \times [0, 1] \setminus \{x\} \times [2^{-n}, 1]$$

- (2) a local base at points of the form $\langle x, y \rangle$, with $y > 0$ consists of the sets

$$U(x, y, n) = \{x\} \times (y - 2^{-n}, y + 2^{-n})$$

We call the resulting space the connected comb and denote it by C . It is straightforward to verify that C is compact, Hausdorff and connected; it is first-countable by definition.

Date: Friday 30-05-2008 at 14:21:32 (cest).

2000 Mathematics Subject Classification. Primary: 54F15. Secondary: 03E50, 03E65, 54A35, 54D40, 54G20.

Key words and phrases. first-countable continuum, continuous image, $\mathbb{H}^*$, Cohen reals.

†Supported by NSF grant DMS-0554896.

For each $x \in [0, 1]$ and positive a we define to be the following cross-shaped closed subset of C^2 :

$$D_{x,a} = (\{x\} \times [a, 1] \times C) \cup (C \times \{x\} \times [a, 1])$$

We note the following two properties of the sets $D_{x,a}$

- (1) if $a < b$ then $D_{x,b}$ is in the interior of $D_{x,a}$, and
- (2) if $x \neq y$ then $D_{x,a} \cap D_{y,a}$ is the union of two squares: $\{x\} \times [a, 1] \times \{y\} \times [a, 1]$ and $\{y\} \times [a, 1] \times \{x\} \times [a, 1]$

Now take any $\aleph_2$ -sized subset of $[0, 1]$ and index it (faithfully) as $\{x_\alpha : \alpha < \omega_2\}$. We use this indexing to identify E with the subset $\{\langle x_\alpha, x_\beta \rangle : \langle \alpha, \beta \rangle \in E\}$ of the unit square. Next we remove from C^2 the following open set:

$$\bigcup_{\langle x, y \rangle \notin E} \left((\{x\} \times (0, 1] \times \{y\} \times (0, 1]) \cup (\{y\} \times (0, 1] \times \{x\} \times (0, 1]) \right)$$

The resulting compact space we denote by C_E . Observe that the intersections $D_{x_\alpha, a} \cap C_E$ represent E in the sense that $D_{x_\alpha, a} \cap D_{x_\beta, a} \cap C_E$ is nonempty if and only if $\langle \alpha, \beta \rangle \in E$. We write $D_{x_\alpha, a}^E = D_{x_\alpha, a} \cap C_E$.

We show that C_E is (arcwise) connected.

To begin: the square S of the base line of C is a subset of C_E and homeomorphic to the unit square so that it is (arcwise) connected.

Let $\langle x, a, y, b \rangle$ be a point of C_E not in S . If, say, $a = 0$ then $\{\langle x, 0 \rangle\} \times (\{y\} \times [0, b])$ is an arc in C_E that connects $\langle x, 0, y, b \rangle$ to the point $\langle x, 0, y, 0 \rangle$ in S . If $a, b > 0$ then $\langle x, y \rangle \in E$ and the whole square $\{x\} \times [0, 1] \times \{y\} \times [0, 1]$ is in C_E and it provides us with an arc in C_E from $\langle x, a, y, b \rangle$ to $\langle x, 0, y, 0 \rangle$.

We find that C_E is a first-countable continuum. It remains to show that it is not an $\mathbb{H}^*$ -image.

Assume $h : \mathbb{H}^* \rightarrow C_E$ is a continuous surjection and consider, for each α , the sets $D_{x_\alpha, \frac{3}{4}}^E$ and $D_{x_\alpha, \frac{1}{2}}^E$.

Using standard properties of $\beta\mathbb{H}$, see [77; 7, Proposition 3.2], we find for each α a sequence $\langle (a_{\alpha, n}, b_{\alpha, n}) : n \in \mathbb{N} \rangle$ of open intervals with rational endpoints, and with $b_{\alpha, n} < a_{\alpha, n+1}$ for all n , such that $h^\leftarrow[D_{x_\alpha, \frac{3}{4}}^E] \subseteq \text{Ex } O_\alpha \cap \mathbb{H}^* \subseteq h^\leftarrow[D_{x_\alpha, \frac{1}{2}}^E]$, where $O_\alpha = \bigcup_n (a_{\alpha, n}, b_{\alpha, n})$. Because the intersections of the sets $D_{x_\alpha, a}^E$ represent E the intersections of the O_α will do this as well: the conditions ' $O_\alpha \cap O_\beta$ is unbounded' and ' $\langle \alpha, \beta \rangle \in E$ ' are equivalent.

In the next subsection we show that for (many) $\langle \alpha, \beta \rangle$ this equivalence does not hold and that therefore C_E is not a continuous image of $\mathbb{H}^*$.

Note also that our continuum is not an $\mathbb{N}^*$ -image either: if $g : \mathbb{N}^* \rightarrow C_E$ were continuous and onto we could use clopen subsets of $\mathbb{N}^*$ and their representing infinite subsets of $\mathbb{N}$ to contradict the unrepresentability property of E .

Destroying the equivalence. We follow the argument from [22, 2] and we rely on Kunen's book [88; 8, Chapter VII] for basic facts on forcing. We let $L = \{\langle \alpha, \beta \rangle \in \omega_2^2 : \alpha \leq \beta\}$ and we force with the partial order $\text{Fn}(L, 2)$ of finite partial functions with domain in L and range in $\{0, 1\}$. If G is a generic filter on $\text{Fn}(L, 2)$ then we let $E = \{\langle \alpha, \beta \rangle : \bigcup G(\alpha, \beta) = 1 \text{ or } \bigcup G(\beta, \alpha) = 1\}$.

To show that E is as required we take a nice name $\dot{F}$ for a function from ω_2 to $(\mathbb{Q}^2)^\omega$ that represents a choice of open sets $\alpha \mapsto O_\alpha$ as in above in that $F(\alpha) = \langle \langle a_{\alpha, n}, b_{\alpha, n} \rangle : n \in \omega \rangle$ for all α . As a nice name $\dot{F}$ is a subset of $\omega_2 \times \omega \times \mathbb{Q}^2 \times \text{Fn}(L, 2)$, where for each point $\langle \alpha, n, a, b \rangle$ the set $\{p : \langle \alpha, n, a, b, p \rangle \in \dot{F}\}$ is a maximal antichain in the set of conditions that forces the n th term of $\dot{F}(\alpha)$ to be $\langle a, b \rangle$.

For each α we let I_α be the set of ordinals that occur in the domains of the conditions that appear as a fifth coordinate in the elements of $\dot{F}$ with first coordinate α . The sets I_α are countable, by the ccc of $\text{Fn}(L, 2)$. We may therefore apply the Free-Set Lemma, see [66; 6, Corollary 44.2], and find a subset A of ω_2 of cardinality $\aleph_2$ such that $\alpha \notin I_\beta$ and $\beta \notin I_\alpha$ whenever $\alpha, \beta \in A$ and $\alpha \neq \beta$.

Let $p \in \text{Fn}(L, 2)$ be arbitrary and take α and β in A with $\alpha < \beta$ and such that $\alpha > \eta$ whenever η occurs in p . Consider the condition $q = p \cup \{\langle \alpha, \beta, 1 \rangle\}$. If q forces $O_\alpha \cap O_\beta$ to be bounded in $[0, \infty)$ then we are done: q forces that the equivalence fails at $\langle \alpha, \beta \rangle$.

If q does not force the intersection to be bounded we can extend q to a condition r that forces $O_\alpha \cap O_\beta$ to be unbounded. We define an automorphism h of $\text{Fn}(L, 2)$ by changing the value of the conditions only at $\langle \alpha, \beta \rangle$: from 0 to 1 and vice versa. The condition p as well as the names $\dot{x}_\alpha$ and $\dot{x}_\beta$ are invariant under h . It follows that $h(r)$ extends p and

$$h(r) \Vdash \bigcup \dot{G}(\alpha, \beta) = 0 \text{ and } O_\alpha \cap O_\beta \text{ is unbounded}$$

so again the equivalence is forced to fail at $\langle \alpha, \beta \rangle$.

Remark. The argument above goes through almost verbatim to show that Bell's graph can also be obtained adding $\aleph_2$ random reals. When forcing with the random real algebra one needs only consider conditions that belong to the σ -algebra generated by the clopen sets of the product $\{0, 1\}^L$; these all have countable supports so that, again by the ccc, one can define the sets I_α as before. The rest of the argument remains virtually unchanged.

REFERENCES

- [1] [cite.MR02516951A. V. Arhangel'skiĭ, *The power of bicomcompacta with first axiom of countability*, Soviet Mathematics Doklady **10** (1969), 951–955. Russian original: Doklady Akademii Nauk SSSR **187** (1969) 967–970. MR 0251695 (40 #4922)
- [2] [cite.MR6778602Murray G. Bell, *The space of complete subgraphs of a graph*, Commentationes Mathematicae Universitatis Carolinae **23** (1982), no. 3, 525–536. MR **677860** (84a:54050)
- [3] [cite.MR10587953_____, *A first countable compact space that is not an N^* image*, Topology and its Applications **35** (1990), no. 2-3, 153–156. MR **1058795** (91m:54028)
- [4] [cite.MR17074894Alan Dow and Klaas Pieter Hart, *A universal continuum of weight $\aleph$* , Transactions of the American Mathematical Society **353** (2001), no. 5, 1819–1838. MR **1707489** (2001g:54037)
- [5] [cite.dowhart-sep5_____, *A separable non-remainder of $\mathbb{H}$* , Proceedings of the American Mathematical Society (to appear), available at <http://arxiv.org/abs/0805.2265>.
- [6] [cite.MR7955926Paul Erdős, András Hajnal, Attila Máté, and Richard Rado, *Combinatorial set theory: partition relations for cardinals*, Studies in Logic and the Foundations of Mathematics, vol. 106, North-Holland Publishing Co., Amsterdam, 1984. MR **795592** (87g:04002)
- [7] [cite.Hart7Klaas Pieter Hart, *The Čech-Stone compactification of the Real Line*, Recent progress in general topology, 1992, pp. 317–352. MR **95g:54004**
- [8] [cite.MR5973428Kenneth Kunen, *Set theory*, Studies in Logic and the Foundations of Mathematics, vol. 102, North-Holland Publishing Co., Amsterdam, 1980. An introduction to independence proofs. MR **597342** (82f:03001)
- [9] [cite.Parovicenko639I. I. Parovičenko, *A universal bicomcompact of weight $\aleph$* , Soviet Mathematics Doklady **4** (1963), 592–595. Russian original: *Ob odnom universal'nom bikompakte vesa $\aleph$* , Doklady Akademii Nauk SSSR **150** (1963) 36–39. MR 27#719

DEPARTMENT OF MATHEMATICS, UNC-CHARLOTTE, 9201 UNIVERSITY CITY BLVD., CHARLOTTE, NC 28223-0001

E-mail address: adow@uncc.edu

URL: <http://www.math.uncc.edu/~adow>

FACULTY OF ELECTRICAL ENGINEERING, MATHEMATICS AND COMPUTER SCIENCE, TU DELFT, POSTBUS 5031, 2600 GA DELFT, THE NETHERLANDS

E-mail address: k.p.hart@tudelft.nl

URL: <http://fa.its.tudelft.nl/~hart>